


[Start](#)
[Author Index](#)
[View Uploaded Presentations](#)
[Meeting Information](#)

Paper No. 200-1

Presentation Time: 8:10 AM

THE CHELYABINSK SUPERBOLIDE: CHARACTERIZATION OF METEOROID AND LESSONS FOR THE EARTH IMPACT HAZARD

KOEBERL, Christian, Department of Lithospheric Research, University of Vienna, Althanstrasse 14, A-1090 Vienna, Austria, also of the Natural History Museum, Burgring 7, A-1010 Vienna, Austria, christian.koeberl@univie.ac.at, FERRIÈRE, Ludovic, Department of Mineralogy and Petrography, Natural History Museum, Burgring 7, Vienna, A-1010, Austria, and BRANDSTAETTER, Franz, Mineralogy and Petrology, Natural History Museum, Burgring 7, Vienna, A1010, Austria

On February 15, 2013, a brilliant fireball (a "superbolide") entered the Earth's atmosphere over Russia at a relatively shallow angle of ca. 18 degrees and exploded at an altitude of at least 23 km (maybe closer to 30 km) at about 9:20 a.m. local time (3:20 a.m. UTC). At the time of the maximum brightness the bolide was travelling at about 18 km/s. The fireball and the explosion were observed over a large area of the southern Ural region, followed by an explosion and shock wave that damaged several buildings and injured more than a thousand people in the region around the city of Chelyabinsk. Thousands of relatively small stones fell about 40 km south of Chelyabinsk and were rapidly recovered, bringing some extremely fresh material for scientific investigations. The largest object recovered so far weighs just a few kg. We recently reported on a multi-disciplinary study, involving mineralogy, petrography, magnetic properties, and cosmogenic radionuclide contents of a dozen stones of the Chelyabinsk meteorite that allowed us to confirm its classification and to characterize the initial meteoroid [1]. Petrographic observations and mineral compositions agree with the classification of the Chelyabinsk meteorite as an LL5 chondrite. Magnetic properties indicate a lack of tetrataenite in the meteorite. The Chelyabinsk meteorite is strongly brecciated and cross-cut by many impact melt veins, which are mainly composed of silicate fragments (unmelted chondritic constituents) surrounded by a matrix containing abundant and finely dispersed sulfides and metal grains.

In total, 12 stones (2-400 g) were measured by [1] for the cosmogenic radionuclides ^{54}Mn ($T_{1/2} = 312$ d), ^{22}Na ($T_{1/2} = 2.60$ y), ^{60}Co ($T_{1/2} = 5.27$ y), and ^{26}Al ($T_{1/2} = 7 \times 10^5$ y). The large variations observed in activities of radionuclides (especially for ^{60}Co) indicate that the samples studied were both from the exterior as well as from the interior of the meteoroid; a pre-explosion meteoroid diameter of 14 to 20 m is indicated [1]. The explosion of this meteoroid, at about 440 kt TNT equivalent, can be compared to the Tunguska event, where no meteorites seem to have fallen, despite a lower explosion altitude. The implications of the two events in terms of danger assessment will be discussed.

[1] Povinec P. et al., 76th Annual Meteoritical Society Meeting (2013), abs. No. 5196.

Session No. 200

T5. Impact Cratering in the Solar System: Fire to Ice—Vacuum to Atmospheres

Tuesday, 29 October 2013: 8:00 AM-12:00 PM

Colorado Convention Center Room 201

 Geological Society of America *Abstracts with Programs*. Vol. 45, No. 7, p.0

© Copyright 2013 The Geological Society of America (GSA), all rights reserved. Permission is hereby granted to the author(s) of this abstract to reproduce and distribute it freely, for noncommercial purposes. Permission is hereby granted to any individual scientist to download a single copy of this electronic file and reproduce up to 20 paper copies for noncommercial purposes advancing science and education, including classroom use, providing all reproductions include the complete content shown here, including the author information. All other forms of reproduction and/or transmittal are prohibited without written permission from GSA Copyright Permissions.

See more of: [Impact Cratering in the Solar System: Fire to Ice—Vacuum to Atmospheres](#)

See more of: [Topical Sessions](#)

<< [Previous Abstract](#) | [Next Abstract](#) >>